

Setup for the next days.

sequence (a_0, a_1, a_2, \dots) $a_n := \#$ ways to build a structure of Type I on $[n]$

sequence (b_0, b_1, b_2, \dots) $b_n := \#$ ways to build a structure of Type II on $[n]$

- Examples for "building structure own"
- choosing an ordering $(n!)$
 - choosing a subset (2^n)
 - choosing an element (n)
 - doing nothing (1)
 - \emptyset (0)


What do the coefficients of the series $C(x) = A(x) \cdot B(x)$ represent? (where $A(x) = \sum_{n=0}^{\infty} a_n x^n$ $B(x) = \sum_{n=0}^{\infty} b_n x^n$)

Proposition $C(x) = A(x) \cdot B(x) := \sum c_n x^n$

(Product Formula)

$\Rightarrow c_n = \#$ of ways to do the following: (maybe empty)

① ~~Build a set~~ Split $[n]$ into two intervals

I_1 and I_2 

② Build structure of Type I on I_1

③ - || - Type II on I_2

PF: ~~Classify~~ Classify according to where do we split: $[i] \cup [n-i]$

$c_n^{(i)} := \#$ possibilities if we split at $i = a_i \cdot b_{n-i}$ (Product Rule)

$\#$ of ways ~~ways~~ $= \sum_{i=0}^n c_n^{(i)}$ (Sum Rule) \Rightarrow ~~ways~~ $\#$ of ways $= \sum_{i=0}^n a_i b_{n-i}$

Also: $(a_0 + a_1 x + a_2 x^2 + \dots) (b_0 + b_1 x + b_2 x^2 + \dots) \Rightarrow \sum_{n=0}^{\infty} (a_0 b_n + a_1 b_{n-1} + \dots + a_n b_0) x^n$

Example: Design term in an Engineering Department
Term: n days

Decisions: How long should theoretical part be?

k days ($1 \leq k \leq n-2$)

Rest is laboratory part
($n-k$ days)

- When should ~~the~~ be the
- x project day for the theoretical part?
- two project days for the lab part?

How many ways are there? $\boxed{f_n}$

Product Formula:

$a_n := \#$ ways to select project day from
an n day long theoretical part

$$a_n = n$$

$b_n := \#$ ways to select TWO project days from
an n day long lab part

$$b_n = \binom{n}{2}$$

$$a(x) = \sum_{n=1}^{\infty} n x^n = x \cdot \left(\sum_{n=0}^{\infty} x^n \right)' = x \cdot \left(\frac{1}{1-x} \right)' = \frac{x}{(1-x)^2}$$


$$b(x) = \sum_{n=2}^{\infty} \binom{n}{2} x^n = \frac{x^2}{2} \cdot \left(\sum_{n=0}^{\infty} x^n \right)'' = \frac{x^2}{2} \cdot \left(\frac{1}{1-x} \right)'' = \frac{x^2}{(1-x)^3}$$

$$f(x) = \sum_{n=3}^{\infty} f_n x^n = a(x) \cdot b(x) = \frac{x^3}{(1-x)^5} = x^3 \sum_{n=0}^{\infty} \binom{n+4}{4} x^n = \sum_{n=3}^{\infty} \binom{n+1}{4} x^n$$

$\Rightarrow \boxed{f_n = \binom{n+1}{4}}$

Catalan numbers

Student



empty jar

every day : either 1 Euro coin in
or 1 Euro coin out

After $2n$ days the jar is empty again.

How many ways can this happen? C_n

Define $c_0 := 1$ Real time exercise c_1, c_2, c_3

$$C_n := \left[\left\{ (b_1, b_2, \dots, b_{2n}) \in (+1, -1)^{2n} : \sum_{j=1}^{2n} b_j = 0 \right. \right. \\ \left. \left. \text{AND } \forall k \sum_{j=1}^k b_j \geq 0 \right\} \right]$$

At the end the jar is empty

The jar never has negative coins
amount of

Classify ~~elements~~ elements according to
~~SMALLEST~~ $k > 0$ when $\sum_{j=1}^k b_j = 0$ (FIRST time when jar is empty again)

of ways to finish afterwards : C_{n-k}

of ways to get there : MUST start with $b_1 = +1$
finish with $b_{2k} = -1$

AND $\sum_{j=2}^k b_j \geq 0 \quad \forall \quad k, 2 \leq k \leq 2n$

There are c_{n-1} of these, $\text{AND } \sum_{j=2}^{2k-1} b_j = 0$

Recurrence: $C_n = \sum_{i=1}^n C_{i-1} C_{n-i} = \sum_{j=0}^{n-1} C_j C_{n-1-j}$

$$C(x) \cdot C(x) = \sum_{l=0}^{\infty} \underbrace{\sum_{i=0}^l C_i C_{l-i}}_{C_{l+1}} x^l = \frac{C(x) - c_0}{x}$$

$$xC(x)^2 - C(x) + 1 = 0$$

$$C(x) = \frac{1 \pm \sqrt{1-4x}}{2x}$$

Is it $C(x) = \frac{1 + \sqrt{1-4x}}{2x}$ or

$C(x) = \frac{1 - \sqrt{1-4x}}{2x}$

CANNOT BE BOTH

when $x \rightarrow 0$
 $\rightarrow \infty$

But it should tend to $c_0 = 1$

So this is the real thing

$$\sqrt{1-4x} = (1-4x)^{\frac{1}{2}} = \sum_{h=0}^{\infty} \binom{\frac{1}{2}}{h} (-4x)^h = 1 + \frac{1}{2}(-4x) + \frac{\frac{1}{2}(-\frac{1}{2})}{2!} (-4)^2 x^2 + \dots$$

$$\dots + \frac{\frac{1}{2}(-\frac{1}{2})(-\frac{3}{2}) \dots (-\frac{1}{2}-h+1)}{h!} (-4)^h x^h + \dots$$

$$= 1 - 2x - 2x^2 - \dots (-1)^{2n-1} \frac{(2n-3)!!}{2^n n!} 4^n x^n$$

$$\frac{1 - \sqrt{1-4x}}{2x} = 1 - \cancel{2x} - \dots + \frac{(2n-3)!! (2^{n-1} (n-1)!!)}{n! (n-1)!} x^{n-1}$$

$$C_n = \frac{1}{n} \frac{(2n-2)!}{(n-1)! (n-1)!} = \frac{1}{n} \binom{2n-2}{n-1}$$

Putting Catalan numbers into the Product Formula setup

Define: • good sequence $(b_1, \dots, b_{2n}) \in \{-1, 1\}^{2n} : \sum_{i=1}^{2n} b_i = 0$

$$\forall j, 1 \leq j \leq 2n \quad \sum_{i=1}^j b_i \geq 0$$

• very good sequence $(v_1, \dots, v_{2n}) \in \{-1, 1\}^{2n} : \sum_{i=1}^{2n} v_i = 0$

$$\forall j, 2 \leq j \leq 2n-1 \quad \sum_{i=1}^j b_i > 0$$

$\forall n \geq 1$ $C_n := \#$ of good sequences of length $2n$

$V_n := \#$ of very good — || —

We have shown $C_n = \sum_{i=1}^n v_i C_{n-i}$ (provided we define $C_0 := 1$)

For Product Formula we need $C_n = \sum_{i=0}^n v_i C_{n-i}$ so define $V_0 = 0$

- Split $[n]$ into intervals I_1, I_2
- Put very good sequence on I_1 (structure of Type I)
- Put good sequence on I_2 (Type II)

\Rightarrow Generating function is $C(x) \cdot V(x)$

It is equal to $C(x)$ EXCEPT $n=0$ ($C_0=1$, but $C_0 \cdot V_0=0$)

$$\text{So } C(x) \cdot V(x) = C(x) - 1$$

Now $V(x) = xC(x)$, since we have shown $V_n = C_{n-1}$
 $\forall n \geq 1$

Indeed, there is a bijection between

good sequences of length $2n-2$ and very good sequences of length $2n$

(v_1, \dots, v_{2n}) very good $\rightarrow (v_2, \dots, v_{2n-1})$ is good

$(+1, b_1, \dots, b_{2n-2}, -1)$ very good $\leftarrow (b_1, \dots, b_{2n-2})$ good

Prop (k-wise Product Formula) For $s=1, 2, \dots, k$, let


$(a_0^{(s)}, a_1^{(s)}, a_2^{(s)}, \dots)$ sequence with

$a_n^{(s)} := \#$ of ways to build structure of type s on $[n]$

~~let~~ let $A_j(x) = \sum_{n=0}^{\infty} a_n^{(j)} x^n$ and let (c_0, c_1, c_2, \dots) be the coefficients in $C(x) = \sum c_n x^n = \prod_{j=1}^k A_j(x)$

Then $c_n = \#$ of ways to do the following:

(1) Split $[n]$ into k intervals $I_1 \cup I_2 \cup \dots \cup I_k$



(2) Build structure of Type j on I_j

~~Pr: ...~~

~~...~~ Induction on k

~~...~~ $k=1 \checkmark$ ($k=2 \checkmark$ deg Product Form)

Do the procedure as follows:

(1) Split $[n]$ into two intervals I^* and I_2

(2) Build the " $(k-1)$ interval" structure with structures of Type $1, 2, \dots, k-1$ on I^*

(3) Build structure of Type k on I_2

~~Generating function of the sequence~~ Generating function of the sequence "How many ways can we do (2) if $|I^*|=n$?"

$$\text{is } \prod_{j=1}^{k-1} A_j(x)$$

Generating function of (3) is $A_k(x) \implies$ By Product Formula $\implies \prod_{j=1}^k A_j(x)$

Composition of Generating Functions

Recall setup: $(a_n), (b_n)$, Structure of Type I, Type II, $A(x), B(x)$

What do the coefficients of $B(A(x))$ represent?

For $B(A(x))$ to make sense, we require $a_0 = 0$

Look at first Special Case when $(b_0, b_1, \dots) = (1, 1, 1, \dots)$
that is when $B(x) = \frac{1}{1-x}$

Prop: $\frac{1}{1-A(x)}$ is the generating fn of the sequence

$\forall n \geq 1$ $h_n := \#$ of ways to do the following

- ① split $[n]$ into nonempty intervals (unspecified number)
- ② build Structure of Type I on each

Define $h_0 := 1$

PF: Classify according to how many nonempty intervals are in the partition: k

$h_n^{(k)} := \#$ of choices with k non-empty intervals in partition

generating function of $h_n^{(k)} : \sum_{n=0}^{\infty} h_n^{(k)} x^n = (A(x))^k$
 \nwarrow k -wise Product Form

Define $h_0^{(2)} := 0$

This is true because $a_0 = 0$, so partitions with empty intervals are not counted, (In the ~~sequence~~ k -wise Product Theorem partitions ~~are~~ with empty intervals were allowed.)

Sum Rule $\Rightarrow h_n = \sum_{z=1}^n h_n^{(z)} = \sum_{z=1}^{\infty} h_n^{(z)}$ since $h_n^{(z)} = 0$ if $z > n$

$\Rightarrow H(x) = \sum_{n=0}^{\infty} h_n x^n = h_0 + \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} h_n^{(k)} x^n = h_0 + \sum_{k=1}^{\infty} \left(\sum_{n=1}^{\infty} h_n^{(k)} x^n \right) = h_0 + \sum_{k=1}^{\infty} (A(x))^k = (A(x))^k$

Example: n soldiers of a squadron stand in line



Sergeant: • splits the line at some places to form smaller units

• names a commander for each unit

$h_n := \#$ of ways this is possible Define $h_0 := 1$

Use Proposition: $\boxed{a_k := k}$ ($\#$ of ways to select commander from $[k]$)
 $\forall k \geq 0$

(Note ~~that~~ $a_0 := 0$)

$$A(x) = \sum_{k=0}^{\infty} kx^k = x \cdot \sum_{k=0}^{\infty} kx^{k-1} = x \left(\sum_{k=0}^{\infty} x^k \right)' = x \left(\frac{1}{1-x} \right)' = \frac{x}{(1-x)^2}$$

$$\Rightarrow \sum_{k=0}^{\infty} h_k x^k = H(x) = \frac{1}{1-A(x)} = \frac{1}{1-\frac{x}{(1-x)^2}} = \frac{(1-x)^2}{1-3x+x^2} = 1 + \frac{x}{1-3x+x^2}$$

$$\frac{x}{1-3x+x^2} = \frac{A}{1-\alpha_1 x} + \frac{B}{1-\alpha_2 x}$$

$$\alpha_1 = \frac{\sqrt{5}+3}{2} \quad \alpha_2 = \frac{-\sqrt{5}+3}{2}$$

$$z^2 - 3z + 1 = 0$$

$$\alpha_{1,2} = \frac{3 \pm \sqrt{5}}{2}$$

$$A+B=0 \text{ and } -\alpha_2 A - \alpha_1 B = 1$$

$$-\alpha_2 A + \alpha_1 A = 1$$

$$B = -A = \frac{1}{\sqrt{5}} \iff A = \frac{1}{\alpha_1 - \alpha_2} = \frac{1}{\sqrt{5}}$$

$$\Rightarrow H(x) = 1 + \frac{1}{\sqrt{5}} \left(\frac{1}{1-\alpha_1 x} - \frac{1}{1-\alpha_2 x} \right) = 1 + \frac{1}{\sqrt{5}} \left(\sum_{n=0}^{\infty} (\alpha_1 x)^n - \sum_{n=0}^{\infty} (\alpha_2 x)^n \right)$$

$$= 1 + \sum_{n=0}^{\infty} \frac{1}{\sqrt{5}} \alpha_1^n x^n - \sum_{n=0}^{\infty} \frac{1}{\sqrt{5}} \alpha_2^n x^n$$

$$\Rightarrow h_n = \frac{1}{\sqrt{5}} \left(\left(\frac{3+\sqrt{5}}{2} \right)^n - \left(\frac{3-\sqrt{5}}{2} \right)^n \right) \text{ and } h_0 = 1$$

Prop (General Composition Formula)

Recall setup: $(a_n), (b_n)$, Structures of Type I, II, $A(x), B(x)$ $a_0 = 0$

$B(A(x))$ is the generating function of the sequence

$g_n :=$ # of ways to do the following

- ① split $[n]$ into non-empty intervals (unspecified number)
- ② build Structure of Type I on each
- ③ build Structure of Type II on the interval of intervals

Pf: $g_0 = b_0$ ✓ ~~For~~ For $n \geq 1$

Again: Classify according to number of nonempty intervals in partition

$h_n^{(k)}$: # of ways to do steps ① and ② with k non-empty intervals

Then $g_n = \sum_{k=0}^{\infty} h_n^{(k)}$ h_k

Define $h_0^{(k)} = 0$

of ways to do step ③, no matter how step ① and ② happened with k intervals ("Product Rule")

~~For~~ $\forall k \geq 1: \sum_{n=0}^{\infty} h_n^{(k)} x^n = (A(x))^k$

Product Formula (again, using $a_0 = 0$)

$$\begin{aligned} \Rightarrow \sum_{n=0}^{\infty} g_n x^n &= g_0 + \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} h_n^{(k)} \cdot b_k x^n = g_0 + \sum_{k=1}^{\infty} b_k \left(\sum_{n=0}^{\infty} h_n^{(k)} x^n \right) \\ &= b_0 + \sum_{k=1}^{\infty} b_k (A(x))^k = B(A(x)) \end{aligned}$$

$(A(x))^k$

Example for full Composition formula

n soldiers in line

- line split at some places forming smaller units
- a subset of the units is chosen for night duty
(possibly empty)

$$a_n = 1 \quad \forall n \quad (\text{No structure on individual units})$$

$$a_0 = 0$$

$$b_n = 2^n \quad (\# \text{ of subsets of } [n])$$

$$A(x) = \sum_{k=1}^{\infty} x^k = \frac{1}{1-x} - 1 = \frac{x}{1-x}$$

$$B(x) = \sum_{k=0}^{\infty} 2^k x^k = \sum_{k=0}^{\infty} (2x)^k = \frac{1}{1-2x}$$

$$B(A(x)) = \frac{1}{1 - \frac{2x}{1-x}} = \frac{1-x}{1-3x} = \frac{1}{1-3x} - \frac{x}{1-3x}$$

$$= \sum_{i=0}^{\infty} (3x)^i - x \sum_{i=0}^{\infty} (3x)^i = 1 + \sum_{i=1}^{\infty} \underbrace{(3^i - 3^{i-1})}_{2 \cdot 3^{i-1}} x^i$$

For $n \geq 1$ soldiers $\Rightarrow \exists 2 \cdot 3^{n-1}$ options

Exponential generating fns

When the sequence (a_0, a_1, a_2, \dots) grows too fast...
 (superexponentially: $a_n \gg K^n \forall K \in \mathbb{R}$)

Def: (a_0, a_1, \dots) sequence of reals

$\hat{A}(x) = \sum_{n=0}^{\infty} a_n \frac{x^n}{n!}$ is the exponential generating

function of the sequence

Example: $a_0 = 1, \forall n \geq 0 \ a_{n+1} = (n+1)a_n - n^2 + 1$

(linear recurrence, but coefficient is NOT constant!)

Closed formula?

$$\begin{aligned} \hat{A}(x) &= \sum_{n=0}^{\infty} a_{n+1} \frac{x^{n+1}}{(n+1)!} = \sum_{n=0}^{\infty} (n+1) \frac{a_n}{(n+1)!} x^{n+1} - \sum_{n=0}^{\infty} \frac{n^2 - 1}{(n+1)!} x^{n+1} \\ &= x \cdot \sum_{n=0}^{\infty} a_n \frac{x^n}{n!} - \sum_{n=1}^{\infty} \frac{x^{n+1}}{(n-1)!} + \sum_{n=0}^{\infty} \frac{x^{n+1}}{n!} \\ &= x \cdot \hat{A}(x) - x^2 e^x + x e^x \end{aligned}$$

$\frac{n-1}{n!} = \frac{1}{(n-1)!} \quad \begin{matrix} n=0 \\ n \geq 1 \end{matrix}$

$$\hat{A}(x)(1-x) = 1 - x^2 e^x + x e^x$$

$$\hat{A}(x) = \frac{1}{1-x} + x e^x = \sum_{n=0}^{\infty} x^n + \sum_{n=1}^{\infty} \frac{x^n}{(n-1)!}$$

$$\Rightarrow \frac{a_n}{n!} = 1 + \frac{1}{(n-1)!} \quad \Rightarrow a_n = n! + n$$

$\forall n \geq 1$

Products of Exponential Generating Functions

$$(a_0, a_1, \dots) \rightsquigarrow \hat{A}(x) = \sum_{n=0}^{\infty} a_n \frac{x^n}{n!}$$

$$(b_0, b_1, \dots) \rightsquigarrow \hat{B}(x) = \sum_{n=0}^{\infty} b_n \frac{x^n}{n!}$$

$$\sum_{n=0}^{\infty} \left(\sum_{k=0}^n \frac{a_k}{k!} \frac{b_{n-k}}{(n-k)!} \right) x^n = \hat{C}(x) = \hat{A}(x) \cdot \hat{B}(x)$$

$$\underbrace{\left(\sum_{k=0}^n \binom{n}{k} a_k b_{n-k} \right)}_{C_n} \frac{x^n}{n!}$$

Then: (Product formula for exponential generating functions)
 $a_n := \#$ of ways to build a structure of Type I on $[n]$
 $b_n :=$ II on II -

$\Rightarrow C_n := \#$ ways to do the following:

- ① Partition $[n]$ into two subsets $A_1, A_2, A_1 \cup A_2 = [n]$
- ② Build structure Type I on A_1
- ③ II on A_2

Then $C_n = \sum_{k=0}^n \binom{n}{k} a_k b_{n-k}$

and $\hat{C}(x) = \hat{A}(x) \hat{B}(x)$

Example: Football coach, n players

- Divide them into two groups A_1, A_2
- Each player in A_1 should take R, W, or G shirt and then form a line
- Players in A_2 should take a B shirt and then form a line

How many ways can this be done? C_n

$$a_k = \underbrace{(3^k)}_{\substack{\# \text{ of ways to} \\ \text{distribute shirts}}} \underbrace{(k!)}_{\substack{\# \text{ of ways to form a line}}}$$

$$b_k = (k!)$$

$$C_n = \sum_{k=0}^n \binom{n}{k} a_k b_{n-k}$$

$$\hat{A}(x) = \sum_{k=0}^{\infty} k! \frac{3^k x^k}{k!} = \sum_{k=0}^{\infty} (3x)^k = \frac{1}{1-3x}$$

$$\hat{B}(x) = \sum_{k=0}^{\infty} k! \frac{x^k}{k!} = \sum_{k=0}^{\infty} x^k = \frac{1}{1-x}$$

$$\hat{C}(x) = \hat{A}(x) \cdot \hat{B}(x) = \frac{1}{1-3x} \cdot \frac{1}{1-x} = \frac{A}{1-3x} + \frac{B}{1-x}$$

$$= \frac{3}{2} \sum_{n=0}^{\infty} (3x)^n - \frac{1}{2} \sum_{n=0}^{\infty} x^n = \sum_{n=0}^{\infty} \left(\frac{3}{2} 3^n - \frac{1}{2} \right) x^n$$

$$\Rightarrow C_n = \frac{3^{n+1} - 1}{2} n!$$

$$\begin{aligned} A + B &= 1 \\ -A - 3B &= 0 \\ B - 1 - 3B &= 0 \\ -1 &= 2B \end{aligned}$$

Derivative of exponential generating fns.

$$\left(\sum_{n=0}^{\infty} a_n \frac{x^n}{n!} \right)' = \sum_{n=1}^{\infty} a_n \frac{x^{n-1}}{(n-1)!} = \sum_{n=0}^{\infty} a_{n+1} \frac{x^n}{n!}$$

Example: Bell numbers $B(n) = \#$ of partitions of $[n]$ (into non-empty parts: $B(n) = \sum_{i=0}^n S(n,i)$)

$$B(n+1) = \sum_{i=0}^n \binom{n}{i} \cdot B(i)$$

Classify according to size $(n-i)$ of the part of element $n+1$

$\binom{n}{i}$ \downarrow
ways to choose the elements NOT in the same part as $n+1$

$B(i)$ \rightarrow # of ways to partition the rest (the elements NOT in the same part as $n+1$)

Product Formula

$$a_i = B(i)$$

$$b_i = 1$$

$$\hat{\alpha}(x) = \sum B(i) \frac{x^i}{i!}$$

$$\hat{\beta}(x) = \sum \frac{x^i}{i!} = e^x$$

$$c_n = \sum_{i=0}^n \binom{n}{i} B(i) = B(n+1)$$

$$\rightarrow \sum_{n=0}^{\infty} c_n \frac{x^n}{n!} = \sum_{n=0}^{\infty} B(n+1) \frac{x^n}{n!} = \left(\sum_{n=0}^{\infty} B(n) \frac{x^n}{n!} \right)'$$

$$\hat{\alpha}(x) e^x = \hat{\alpha}(x), \hat{\beta}(x) = \hat{\alpha}(x)'$$

$$e^x = \frac{\hat{\alpha}(x)'}{\hat{\alpha}(x)} = (\ln \hat{\alpha}(x))'$$

$$e^x + C = \ln \hat{\alpha}(x)$$

$$x=0 \rightarrow \ln \hat{\alpha}(0) = 0 \Rightarrow C = -1$$

$$e^{e^x - 1} = \hat{\alpha}(x)$$

Prop (k -wise Product Formula for Exponential Generating Functions)

~~Defn~~ For $j=1, 2, \dots, k$, let $(a_0^{(j)}, a_1^{(j)}, \dots)$ sequence such that $a_n^{(j)} := \#$ of ways to build structure of Type j on $[n]$.

$$\text{Let } \hat{A}_j(x) := \sum_{n=0}^{\infty} a_n^{(j)} \frac{x^n}{n!}$$

Then $\prod_{j=1}^k \hat{A}_j(x)$ is the exponential generating function for the sequence

$d_n := \#$ of ways to do the following (maybe empty)

- ① ~~Partition~~ Partition $[n]$ into k subsets S_1, S_2, \dots, S_k
- ② Build structure of Type j on S_j

Pf: Induction on k

• $k=1, 2$ ✓

• Do the procedure as follows:

- ① Partition $[n]$ into two subsets S^* and S_k
- ② Build the " $(k-1)$ -subset"-structure on S^*
- ③ Build Structure ~~on~~ ^{Type} k on S_k

~~Generating~~ Exp. Generating function for the counting sequence of the procedure in ② is $\prod_{j=1}^{k-1} \hat{A}_j(x)$ (by induction)

Exponential generating function for the counting sequence in ③ is $\hat{A}_k(x)$

So by the (pairwise) Product Formula, the exponential generating function for the whole process is

$$\prod_{j=1}^k \hat{A}_j(x)$$

Composition of Exponential Generating Functions

Recall setup: $(a_n), (b_n)$, Structure of Type I, II, $\hat{A}(x), \hat{B}(x)$

What do the coefficients of $\hat{B}(\hat{A}(x))$ represent?

Again to make sense to $\hat{B}(\hat{A}(x))$ we require $\boxed{a_0=0}$

Special Case: $b_n = 1 \forall n \geq 0$, that is $\hat{B}(x) = e^x$

Prop: $e^{A(x)}$ is the e.g.f. of the sequence

$h_n := \#$ of ways to do the following:

- ① Partition $[n]$ into nonempty subsets ~~(unspecified number)~~ (unspecified number)

Define $\boxed{h_0=0}$ ② Build a structure of Type I on each

Pf: Classify according the number of nonempty subsets in the partition

$h_n^{(k)} := \#$ of ~~choices~~ partitions with k non-empty subsets in partition

Then \bullet Sum Rule $\Rightarrow \boxed{h_n = \sum_{k=1}^n h_n^{(k)} = \sum_{k=1}^{\infty} h_n^{(k)}$ since $h_n^{(k)} = 0 \forall k > n$

\bullet Exponential Generating function: $\sum_{k=0}^{\infty} h_n^{(k)} \frac{x^k}{k!} = \frac{1}{k!} (\hat{A}(x))^k$

① Product Formula also counts partitions with ^{some} empty sets; these have no contribution here, since $a_0=0$ ✓
 ② Product Formula must be divided by $k!$, since ~~partitions with empty sets~~ are not here sets of partition ~~are not~~ labels

$$\hat{H}(x) = \sum_{n=0}^{\infty} h_n \frac{x^n}{n!} = h_0 + \sum_{n=1}^{\infty} \sum_{k=1}^{\infty} h_n^{(k)} \frac{x^n}{n!} = 1 + \sum_{k=1}^{\infty} \left(\sum_{n=1}^{\infty} h_n^{(k)} \frac{x^n}{n!} \right) \frac{1}{k!} = e^{\hat{A}(x)}$$

Example: Arrange n people into groups and seat them around circular table.

Sol: $a_k = (k-1)! \quad \forall k \geq 1$
 $a_0 = 0$
 $b_k = 1 \quad \forall k \geq 0$

$$\hat{A}(x) = \sum_{k=0}^{\infty} (k-1)! \frac{x^k}{k!} = \sum_{k=1}^{\infty} \frac{x^k}{k} = \ln \frac{1}{1-x}$$

$$\Rightarrow \hat{G}(x) = e^{\ln(\frac{1}{1-x})} = \frac{1}{1-x} = \sum_{k=0}^{\infty} x^k = \sum_{k=0}^{\infty} \frac{k!}{k!} \frac{x^k}{k!}$$

Real Time Exercise: Combinatorial solution?

Example: # of partition of $[n]$ into block of sizes 3, 4, or 9

\hat{f}_n

$$\hat{F}(x) := \sum_{k=0}^{\infty} f_k \frac{x^k}{k!}$$

$$a_k := \begin{cases} 1 & k=3 \\ 0 & k \neq 3 \end{cases}$$

$$b_k := \begin{cases} 1 & k=4 \\ 0 & k \neq 4 \end{cases}$$

$$c_k := \begin{cases} 1 & k=9 \\ 0 & k \neq 9 \end{cases}$$

$$\hat{A}(x) = \frac{x^3}{3!}$$

$$\hat{B}(x) = \frac{x^4}{4!}$$

$$\hat{C}(x) = \frac{x^9}{9!}$$

$e^{\hat{A}(x)}$ e.g.f. of

of partitioning into parts of size 3, (choice of a_k makes sure that in a partition of $[n]$ into subsets only those are counted where each part is of size 3)

$e^{\hat{B}(x)}$

$e^{\hat{C}(x)}$

$e^{\hat{A}(x) + \hat{B}(x) + \hat{C}(x)}$ is e.g.f. of partitioning into three parts: first to be partitioned into 3s, second to be partitioned into 4s, third to be partitioned into 9s

Composition of Exponential Generating Fns

~~The Composition Formula~~

~~##~~

$$(a_0, a_1, \dots) \rightsquigarrow \hat{A}(x) = \sum_{n=0}^{\infty} a_n \frac{x^n}{n!}$$

$$(b_0, b_1, \dots) \rightsquigarrow \hat{B}(x) = \sum_{n=0}^{\infty} b_n \frac{x^n}{n!}$$

$$\sum_{k=0}^{\infty} \underbrace{b_k \frac{\hat{A}(x)^k}{k!}}_{\text{Makes sense only if } a_0 = 0} = \hat{B}(\hat{A}(x))$$

Makes sense only if $a_0 = 0$

Then: (Composition Formula of Exponential Generating Fns)

$a_n :=$ # of ways to build a structure of Type I on $[n]$

$b_n :=$ # " " " " Type II on $[n]$

$c_n :=$ # ways to do the following:

- ① Partition $[n]$ into nonempty subsets (unspecified number)
- ② Build structure of Type I on each ~~set~~ subset
- ③ Build " " " " Type II on set of subsets

$$\hat{C}(x) = \sum c_n \frac{x^n}{n!} = \hat{B}(\hat{A}(x))$$

\mathcal{P} : classify according to # of parts in partition:

$c_n^{(k)} =$ # of ways to do above with k parts (nonempty) $\Rightarrow c_n = \sum_{k=1}^n c_n^{(k)}$

$$\hat{C} = \sum_{k=0}^{\infty} \hat{C}^{(k)}(x)$$

$$\hat{C}^{(k)}(x) = \underbrace{b_k}_{\text{put structure of Type II on } [k]} \cdot \underbrace{\hat{A}^k(x)}_{\text{Product formula for } k \text{ parts}} \cdot \frac{1}{k!}$$

put structure of Type II on $[k]$ Product formula for k parts

Example: n distinct cards

- Split them into (non-empty) decks, each with an even number of cards
- order each deck
- order decks in a line

How many ways can we do this? g_n

Composition Formula:

$$a_n := \begin{cases} n! & n \text{ is even } \geq 2 \\ 0 & n \text{ is odd or } n=0 \end{cases}$$

$$b_n := n! \quad \forall n \geq 0$$

$$\hat{A}(x) = \sum_{n=0}^{\infty} a_n \frac{x^n}{n!} = \sum_{\substack{n \geq 2 \\ n \text{ is even}}} x^n = \frac{x^2}{1-x^2}$$

$$\hat{B}(x) = \sum_{n=0}^{\infty} b_n \frac{x^n}{n!} = \sum_{n=0}^{\infty} x^n = \frac{1}{1-x}$$

$$\begin{aligned} \Rightarrow \hat{G}(x) &= \hat{B}(\hat{A}(x)) = \frac{1}{1 - \frac{x^2}{1-x^2}} = \frac{1-x^2}{1-2x^2} \\ &= 1 + \frac{x^2}{1-2x^2} = 1 + x^2 \sum_{m=0}^{\infty} (2x^2)^m = 1 + \sum_{m=0}^{\infty} 2^m x^{2m+2} \end{aligned}$$

~~$\Rightarrow g_n = \dots$~~

$$g_{2m+1} = 0 \quad \forall m$$
$$g_{2m} = 2^{m-1} (2m)! \quad \forall m$$

$$\Rightarrow g_n = 2^{\frac{n}{2}-1} \cdot n!$$